

STATE/OPERATOR CORRESPONDENCE IN HIGHER-SPIN DS/CFT

Gim Seng Ng and Andrew Strominger

*Center for the Fundamental Laws of Nature, Harvard University,
Cambridge, MA 02138, USA*

Abstract

A recently conjectured microscopic realization of the dS_4/CFT_3 correspondence relating Vasiliev's higher-spin gravity on dS_4 to a Euclidean $Sp(N)$ CFT_3 is used to illuminate some previously inaccessible aspects of the dS/CFT dictionary. In particular it is argued that states of the boundary CFT_3 on S^2 are holographically dual to bulk states on geodesically complete, spacelike R^3 slices which terminate on an S^2 at future infinity. The dictionary is described in detail for the case of free scalar excitations. The ground states of the free or critical $Sp(N)$ model are dual to dS -invariant plane-wave type vacua, while the bulk Euclidean vacuum is dual to a certain mixed state in the CFT_3 . CFT_3 states created by operator insertions are found to be dual to (anti) quasinormal modes in the bulk. A norm is defined on the R^3 bulk Hilbert space and shown for the scalar case to be equivalent to both the Zamolodchikov and pseudounitary C-norm of the $Sp(N)$ CFT_3 .

Contents

1	Introduction	1
2	Global dS vacua at $m^2\ell^2 = 2$	4
2.1	Modes	5
2.1.1	Neumann and Dirichlet modes	5
2.1.2	Euclidean modes	6
2.2	Vacua	7
3	Boundary operators and quasinormal modes	10
3.1	Highest and lowest weight wavefunctions	10
3.2	General multi-operator insertions	13
3.3	Klein-Gordon inner product	14
4	The southern Hilbert space	15
4.1	States	15
4.2	Norm	17
5	Boundary dual of the bulk Euclidean vacuum	19
6	Pseudounitariness and the C-norm in the $Sp(N)$ CFT₃	20
A	Appendix: dS₄ Killing vectors	22

1 Introduction

The conjectured dS/CFT correspondence attempts to adapt the wonderful successes of the AdS/CFT correspondence to universes (possibly like our own) which exponentially expand in the far future. The hope [1, 2, 3, 4, 5] is to define bulk de Sitter (dS) quantum gravity in terms of a holographically dual CFT living at $\mathcal{I}^+$ of dS, which is the asymptotic conformal boundary at future null infinity. A major obstacle to this program has been the absence of any explicit microscopic realization. This has so far prevented the detailed development of the dS/CFT dictionary. This situation has recently been improved by an explicit proposal [6] relating Vasiliev’s higher-spin gravity in dS₄ [7, 8] to the dual $Sp(N)$ CFT₃ described in [9]. In this paper we will use this higher-spin context to write some new entries in the dS/CFT dictionary.

The recent proposal [6] for a microscopic realization of dS/CFT begins with the duality relating the free (critical) $O(N)$ CFT₃ to higher-spin gravity on AdS₄ with Neumann (Dirichlet) boundary conditions on the scalar field. Higher-spin gravity - unlike string theory [2] - has a simple analytic continuation from negative to positive cosmological constant Λ . Under this continuation, AdS₄ $\rightarrow$ dS₄ and the (singlet) boundary CFT₃ correlators are simply transformed by the replacement of $N \rightarrow -N$. These same transformed correlators arise from the $Sp(N)$ models constructed from anticommuting scalars. It follows that the free (critical) $Sp(N)$ correlators equal those of higher-spin gravity on dS₄ with future Neumann (Dirichlet) scalar boundary conditions (of the type described in [10]) at $\mathcal{I}^+$.

This mathematical relation between the bulk dS and boundary $Sp(N)$ correlators may provide a good starting point for understanding quantum gravity on dS, but so far important physical questions remain unanswered. For example we do not know how to relate these physically *unmeasurable* correlators to a set of true physical observables or to the dS horizon entropy. These crucial entries in the dS/CFT dictionary are yet to be written.

As a step in this direction, in this paper we investigate the relation between quantum states in the bulk higher-spin gravity and those in the boundary CFT₃. Bulk higher-spin gravity has fields of Φ^s with all even spins $s = 0, 2, \dots$, which are dual to CFT₃ operators $\mathcal{O}^s$ with the same spins. In the CFT₃, we can also associate a state to each operator by the state-operator correspondence. One way to do this is to take the southern hemisphere of S^3 , insert the operator $\mathcal{O}^s$ at the south pole, and then define a state $\Psi_{S^2}^s$ as a functional of the boundary conditions on the equatorial S^2 . For every object in the CFT₃, we expect a holographically dual object in the bulk dS₄ theory. This raises the question: what is the bulk representation of the spin- s state $\Psi_{S^2}^s$?

In Lorentzian AdS₄ holography, the state created by a primary operator $\mathcal{O}$ in the CFT₃ on S^2 has, at weak coupling, a bulk representation as the single particle state of the field Φ dual to $\mathcal{O}$ with a smooth minimal-energy wavefunction localized near the center of AdS₄. The form of the wavefunction is dictated by the conformal symmetry.

In dS₄ holography, the situation is rather different. States in dS₄ quantum gravity are usually thought of as wavefunctions on complete spacelike slices which are topologically S^3 .¹ These do not seem to be good candidates for bulk duals to $\Psi_{S^2}^s$ because, among other reasons, they are not associated to any S^2 in $\mathcal{I}^+$. However, dS₄ also has everywhere spacelike and geodesically complete R^3 slices which end at an S^2 in $\mathcal{I}^+$. Here we propose a construction of

¹As explained in [4, 5], such states do play an important role in dS/CFT, but as generating functions for correlators rather than as duals to CFT₃ states on S^2 . The relation between the R^3 and S^3 bulk states in our example is detailed below.

the bulk version of $\Psi_{S^2}^s$ on these slices, again as single particle states whose form is dictated by the conformal symmetry. Interestingly, the classical wavefunction for the particle turns out to be the (anti) quasinormal modes for the static patch of de Sitter, as constructed in [11, 12].²

This relation between bulk and boundary states has a potentially profound nonperturbative consequence briefly mentioned in section 4.1 [14]. The operator $\mathcal{O}^0$ dual to a scalar Φ is bilinear in boundary fermions and hence obeys $(\mathcal{O}^0)^{\frac{N}{2}+1} = 0$. Under bulk-boundary duality this translates into an $\frac{N}{2}$ -adicity relation for Φ : one cannot put more than $\frac{N}{2}$ bulk scalar quanta into the associated quasinormal mode. Further investigation of this dS exclusion principle is deferred to later work.

We also construct a norm for these bulk states and show that it is the Zamolodchikov norm on S^3 of the CFT_3 operator $\mathcal{O}^s$. Explicit formulae are exhibited only for the scalar $s = 0$ case but we expect the construction to generalize to all s .

This paper is organized as follows. In section 2 we revisit the issue of the usual global dS-invariant vacua for a free massive scalar field, paying particular attention to the case of $m^2\ell^2 = 2$ (where ℓ is the de Sitter radius) arising in higher spin gravity. The invariant vacua include the familiar Bunch-Davies Euclidean vacuum $|0_E\rangle$, as well as a pair of $|0^\pm\rangle$ of in/out vacua with no particle production. As the scalar field acting on $|0^-\rangle$ ($|0^+\rangle$) obeys Dirichlet (Neumann) boundary conditions on $\mathcal{I}^+$, these are related to the critical (free) $Sp(N)$ model. Generically all dS-invariant vacua are Bogolyubov transformations of one another, but we find that at $m^2\ell^2 = 2$ the transformation is singular and the in/out vacua are non-normalizable plane-wave type states. In section 3 we use the conformal symmetries to find the classical bulk wavefunctions associated to an operator insertion on $\mathcal{I}^+$, and note the relation to (anti) quasinormal modes. The construction uses a rescaled bulk-to-boundary Green function defined with Neumann or Dirichlet $\mathcal{I}^+$ boundary conditions. We also show that the Klein-Gordon inner product of these wavefunctions agrees with the conformally-covariant CFT_3 operator two-point function. In section 4 we consider the Hilbert space on R^3 slices ending on an S^2 on $\mathcal{I}^+$. This Hilbert space was explicitly constructed in [15] for a free scalar on hyperbolic slices ending on $\mathcal{I}^+$. There are two such Hilbert spaces, which we denote the northern and southern Hilbert space, which live on spatial R^3 slices extending to the north or south of the S^2 . The northern and southern slices add up to a global S^3 . Hence the tensor product of the northern and southern Hilbert spaces is the global Hilbert space on S^3 , much as the left and right Rindler Hilbert spaces tensor to the global

²We are grateful to D. Anninos for pointing this out [13].

Minkowski Hilbert space. We show that the global $|0^\pm\rangle$ vacua are simple tensor products of the northern and southern Dirichlet and Neumann vacua. We then use symmetries to uniquely identify the states of the southern Hilbert space with those of the free and critical $Sp(N)$ models on an S^2 . This leads directly to the dS exclusion principle. We further construct an inner product for the southern Hilbert space which agrees, for states dual to $\mathcal{I}^+$ operator insertions, to the conformal two-point function on $\mathcal{I}^+$. In section 5 we discuss the restriction of Euclidean vacuum to a southern state and recall from [15], that this is a mixed state which is thermal with respect to an $SO(3,1)$ Casimir. It would be interesting to relate this result to dS entropy in the present context. In section 6 we show that the standard CFT_3 state-operator correspondence maps the known pseudo-unitary C-norm of the $Sp(N)$ model to the Zamolodchikov two-point function. This completes the demonstration that the bulk states on R_S^3 have the requisite properties to be dual to the boundary $Sp(N)$ CFT_3 states on S^2 . Speculations are made on the possible relevance of pseudo-unitarity to the consistency of dS/CFT in general. An appendix gives some explicit formulae for the $SO(4,1)$ Killing vectors of dS_4 .

2 Global dS vacua at $m^2\ell^2 = 2$

In this section we describe the quantum theory of a free scalar field Φ in dS_4 with wave equation

$$(\nabla^2 - m^2)\Phi = 0, \tag{2.1}$$

and mass

$$m^2\ell^2 = 2. \tag{2.2}$$

This is the case of interest for Vasiliev's higher-spin gravity. While there have been many general discussions of this problem, peculiar singular behavior as well as simplifications appear at the critical value $m^2\ell^2 = 2$ which are highly relevant to the structure of dS/CFT. A parallel discussion of de Sitter vacua and scalar Green functions in the context of dS/CFT was given in [16]. However that paper in many places specialized to the large mass regime $m^2\ell^2 > \frac{9}{4}$, excluding the region of current interest. The behavior in the region $m^2\ell^2 < \frac{9}{4}$ divides into three cases $m^2\ell^2 > 2$, $m^2\ell^2 = 2$ and $m^2\ell^2 < 2$. Much of the structure we describe below pertains to the entire range $m^2\ell^2 < \frac{9}{4}$ with an additional branch-cut prescription for the Green functions.

2.1 Modes

We will work in the dS_4 global coordinates

$$\frac{ds^2}{\ell^2} = -dt^2 + \cosh^2 t d\Omega_3^2 = -dt^2 + \cosh^2 t [d\psi^2 + \sin^2 \psi (d\theta^2 + \sin^2 \theta d\phi^2)], \quad (2.3)$$

where $\Omega^i \sim (\psi, \theta, \phi)$ are coordinates on the global S^3 slices. Following the notation of [16] solutions of the wave equation can be expanded in modes

$$\phi_{Lj}(x) = y_L(t) Y_{Lj}(\Omega) \quad (2.4)$$

of total angular momentum L and spin labeled by the multi-index j . The spherical harmonics Y_{Lj} obey

$$\begin{aligned} Y_{Lj}^*(\Omega) &= (-)^L Y_{Lj}(\Omega) = Y_{Lj}(\Omega_A), \\ D^2 Y_{Lj}(\Omega) &= -L(L+2) Y_{Lj}(\Omega), \\ \int_{S^3} \sqrt{h} d^3\Omega Y_{Lj}^*(\Omega) Y_{L'j'}(\Omega) &= \delta_{L,L'} \delta_{j,j'}, \\ \sum Y_{Lj}^*(\Omega) Y_{Lj}(\Omega') &= \frac{1}{\sqrt{h}} \delta^3(\Omega - \Omega'), \end{aligned} \quad (2.5)$$

where $\sqrt{h}$ and D^2 are the measure and Laplacian on the unit S^3 , Ω_A is the antipodal point of Ω , and here and hereafter $\sum$ denotes summation over all allowed values of L and j . The time dependence is then governed by the second order ODE

$$\partial_t^2 y_L + 3 \tanh t \partial_t y_L + \left(m^2 \ell^2 + \frac{L(L+2)}{\cosh^2 t} \right) y_L = 0. \quad (2.6)$$

2.1.1 Neumann and Dirichlet modes

Eq. (2.6) has the *real* solutions

$$y_L^\pm = 2^{L+h_\pm+\frac{1}{2}} (L+1)^{\pm\frac{1}{2}} \cosh^L t e^{-(L+h_\pm)t} F\left(L+\frac{3}{2}, L+h_\pm, h_\pm-\frac{1}{2}; -e^{-2t}\right) \quad (2.7)$$

where

$$h_\pm \equiv \frac{3}{2} \pm \sqrt{\frac{9}{4} - m^2 \ell^2}. \quad (2.8)$$

We are interested in $m^2 \ell^2 = 2$, which implies

$$h_- = 1, \quad h_+ = 2, \quad (2.9)$$

and

$$y_L^\pm = \frac{(-i)^{\frac{1}{2} \pm \frac{1}{2}} 2^L}{\sqrt{1+L}} \cosh^L t e^{-(L+1)t} \left[\frac{1}{(1 - ie^{-t})^{2L+2}} \mp \frac{1}{(1 + ie^{-t})^{2L+2}} \right]. \quad (2.10)$$

The modes behave near $\mathcal{I}^+$ as $e^{-h_\pm t}$

$$\begin{aligned} t &\rightarrow \infty, \\ y_L^- &\rightarrow (2(L+1)^{-\frac{1}{2}})e^{-t} + \mathcal{O}(e^{-3t}) && \text{Neumann,} \\ y_L^+ &\rightarrow (4(L+1)^{\frac{1}{2}})e^{-2t} + \mathcal{O}(e^{-4t}) && \text{Dirichlet.} \end{aligned} \quad (2.11)$$

Accordingly we refer to the $+$ modes as Dirichlet and the $-$ modes as Neumann. We have normalized so that the Klein-Gordon inner product is

$$\langle \phi_{Lj}^+ | \phi_{L'j'}^- \rangle_{S^3} \equiv i \int_{S^3} d^3 \Sigma^\mu \phi_{Lj}^{+*} \overleftrightarrow{\partial}_\mu \phi_{L'j'}^- = i \delta_{LL'} \delta_{jj'}, \quad (2.12)$$

with $d^3 \Sigma^\mu$ the induced measure times the normal to the S^3 slice.

Under time reversal

$$y_L^\pm(t) = \pm (-)^L y_L^\pm(-t), \quad (2.13)$$

so that

$$\phi_{Lj}^\pm(x) = \pm \phi_{Lj}^\pm(x_A) = (-)^L \phi_{Lj}^{\pm*}(x), \quad (2.14)$$

where the point x_A is antipodal to the point x . This implies that an incoming Dirichlet (Neumann) mode propagates to an outgoing Dirichlet (Neumann) mode. This is not the case for generic m^2 and, as will be seen below, allows for Dirichlet and Neumann vacua with no particle production.

2.1.2 Euclidean modes

Euclidean modes are defined by the condition that when dS_4 is analytically continued to S^4 they remain nonsingular on the southern hemisphere. In other words

$$y_L^E(t = -\frac{i\pi}{2}) = \text{nonsingular}. \quad (2.15)$$

One finds that the combination

$$y_L^E = \frac{y_L^- + i y_L^+}{\sqrt{2}} = \frac{2^{L+1}}{\sqrt{2L+2}} \frac{\cosh^L t e^{-(L+1)t}}{(1 - ie^{-t})^{2L+2}} \quad (2.16)$$

is nonsingular at $t = -i\pi/2$. Hence $y_L^\mp$ are simply the real and imaginary parts of the y_L^E . (2.13) and (2.14) imply the relations

$$y_L^{E*}(t) = (-)^{L+1} y_L^E(-t) \quad (2.17)$$

$$\phi_{Lj}^E(x_A) = (-)^{L+1} \phi_{Lj}^{E*}(x). \quad (2.18)$$

$$\langle \phi_{Lj}^E | \phi_{L'j'}^E \rangle_{S^3} = \delta_{LL'} \delta_{jj'} \quad (2.19)$$

2.2 Vacua

In the quantum theory Φ is promoted to an operator which we denote $\hat{\Phi}$ obeying the equal time commutation relation

$$[\hat{\Phi}(\Omega, t), \partial_t \hat{\Phi}(\Omega', t)] = \frac{i}{\sqrt{h} \cosh^3 t} \delta^3(\Omega - \Omega'). \quad (2.20)$$

Defining annihilation and creation operators

$$a_{Lj}^E = \langle \phi_{Lj}^E | \hat{\Phi} \rangle_{S^3}, \quad a_{Lj}^{E\dagger} = -\langle \phi_{Lj}^{E*} | \hat{\Phi} \rangle_{S^3}, \quad (2.21)$$

the global Euclidean (or Bunch-Davies) vacuum is defined by

$$a_{Lj}^E |0_E\rangle = 0. \quad (2.22)$$

We normalize so that $\langle 0_E | 0_E \rangle = 1$. For any m^2 there is a family of dS-invariant vacua labeled by a complex parameter α . They are annihilated by the normalized Bogolyubov-transformed oscillators

$$a_{Lj}^\alpha = \frac{1}{\sqrt{1 - e^{\alpha + \alpha^*}}} \left(a_{Lj}^E - e^{\alpha^*} a_{Lj}^{E\dagger} \right). \quad (2.23)$$

We are interested in the vacua annihilated by the Dirichlet or Neumann modes for the case of $m^2 \ell^2 = 2$, which correspond to $e^\alpha = \pm 1$. In that case the Bogolyubov transformation is singular. Nevertheless we can still construct non-normalizable plane-wave type vacua as follows.

The field operator may be decomposed as

$$\hat{\Phi} = \hat{\Phi}^+ + \hat{\Phi}^-, \quad (2.24)$$

where $\hat{\Phi}^\pm \sim e^{-h_\pm t}$ near $\mathcal{I}^+$. The squeezed states

$$|0^\pm\rangle = e^{\pm\frac{1}{2}\Sigma(-)^L(a_{Lj}^{E\dagger})^2}|0_E\rangle \quad (2.25)$$

then obey

$$\hat{\Phi}^-|0^-\rangle = 0 \quad \text{Dirichlet}, \quad (2.26)$$

$$\hat{\Phi}^+|0^+\rangle = 0 \quad \text{Neumann}. \quad (2.27)$$

Since only Dirichlet (Neumann) modes act non-trivially on $|0^-\rangle$ ($|0^+\rangle$) we refer to it as the Dirichlet (Neumann) vacuum. These vacua are dS invariant. With the conventional norm, $\hat{\Phi}^\pm$ are hermitian and their eigenstates are non-normalizable. Generalized dS non-invariant plane-wave type Neumann states with nonzero eigenvalues for $\hat{\Phi}^+$

$$\hat{\Phi}^+|\Phi^+\rangle = \Phi^+|\Phi^+\rangle \quad (2.28)$$

are constructed as

$$|\Phi^+\rangle = e^{-\langle\Phi^+|\hat{\Phi}^-\rangle_{S^3}}|0^+\rangle. \quad (2.29)$$

Φ^+ here is an arbitrary solution of the classical wave equation, which can be parameterized by an arbitrary function $\Phi^+(\Omega)$ on $\mathcal{I}^+$

$$t \rightarrow \infty, \quad \Phi^+(\Omega, t) \rightarrow \Phi^+(\Omega)e^{-h_+t}. \quad (2.30)$$

The states are delta-functional normalizable with respect the usual inner product

$$\langle\Phi^+|\Phi^{+\prime}\rangle = \delta\left(\Phi^+ - \Phi^{+\prime}\right), \quad (2.31)$$

where the delta function integrates to one with the measure

$$\mathcal{D}\Phi^+ \equiv \prod_{L,j} \frac{dc_{Lj}^+}{\sqrt{\pi}}, \quad \Phi^+(x) = \sum c_{Lj}^+ \phi_{Lj}^+(x). \quad (2.32)$$

The c_{Lj}^+ satisfies the reality condition $c_{Lj}^{+*} = c_{Lj}^+(-)^L$. One may similarly define generalized Dirichlet states obeying

$$\hat{\Phi}^-|\Phi^-\rangle = \Phi^-|\Phi^-\rangle. \quad (2.33)$$

The Euclidean vacuum can be expressed in terms of $|0^\pm\rangle$ as

$$|0_E\rangle = \int \mathcal{D}\Phi^\pm e^{\mp \frac{1}{16} \int d^3\Omega d^3\Omega' \Phi^\pm(\Omega) \Delta_\mp(\Omega, \Omega') \Phi^\pm(\Omega')} |\Phi^\pm\rangle, \quad (2.34)$$

where

$$\begin{aligned} \Delta_\pm(\Omega, \Omega') &= \mp \sum Y_{Lj}^*(\Omega) Y_{Lj}(\Omega') (2L+2)^{\pm 1} = \frac{1}{2^{2\mp 1} \pi^2} \frac{1}{(1 - \cos \Theta_3)^{h_\pm}}, \\ \cos \Theta_3(\Omega, \Omega') &\equiv \cos \psi \cos \psi' + \sin \psi \sin \psi' (\cos \theta \cos \theta' + \sin \theta \sin \theta' \cos(\phi - \phi')). \end{aligned} \quad (2.35)$$

$\Delta_\pm$ are the (everywhere positive) two-point functions for a CFT₃ operator with $h_+ = 2$ and $h_- = 1$.³ These satisfy

$$- \int \sqrt{h} d^3\Omega'' \Delta_+(\Omega, \Omega'') \Delta_-(\Omega'', \Omega') = \frac{1}{\sqrt{h}} \delta^3(\Omega - \Omega'). \quad (2.36)$$

We also have the relations

$$|\Phi^+\rangle = \frac{1}{N_0} \int \mathcal{D}\Phi^- e^{\langle \Phi^- | \Phi^+ \rangle_{S^3}} |\Phi^-\rangle, \quad N_0 \equiv \prod_{L,j} \sqrt{2}, \quad (2.37)$$

$$\langle \Phi^- | \Phi^+ \rangle_{S^3} = \frac{1}{N_0} e^{\langle \Phi^- | \Phi^+ \rangle_{S^3}}. \quad (2.38)$$

In particular

$$\langle 0^- | 0^+ \rangle_{S^3} = \frac{1}{N_0}. \quad (2.39)$$

The Wightman function in the Euclidean vacuum is

$$\begin{aligned} G_E(x; x') &= \sum \phi_{Lj}^E(x) \phi_{Lj}^{E*}(x') \\ &= \sum (-)^{L+1} \phi_{Lj}^E(x) \phi_{Lj}^E(x'_A) \\ &= \frac{1}{2} \sum (-)^L (\phi_{Lj}^-(x) \phi_{Lj}^-(x') + \phi_{Lj}^+(x) \phi_{Lj}^+(x') + i \phi_{Lj}^+(x) \phi_{Lj}^-(x') - i \phi_{Lj}^-(x) \phi_{Lj}^+(x')). \end{aligned} \quad (2.40)$$

In terms of the dS-invariant distance function

$$P(t, \Omega; t', \Omega') = \cosh t \cosh t' \cos \Theta_3(\Omega, \Omega') - \sinh t \sinh t', \quad (2.41)$$

³Here, we regulate the expressions of $\Delta_\pm$ as sums over the spherical harmonics by introducing $e^{-L\varepsilon}$ in each term in the sum and take the limit of $\varepsilon \rightarrow 0$ at the end after the summation.

this becomes simply

$$G_E(x; x') = \frac{1}{8\pi^2} \frac{1}{1 - P(x; x')}, \quad (2.42)$$

with the usual $i\varepsilon$ prescription for the singularity.

3 Boundary operators and quasinormal modes

According to the dS_4/CFT_3 dictionary, for every spin zero primary CFT_3 operator $\mathcal{O}$ of conformal weight h there is a bulk scalar field Φ with mass $m^2\ell^2 = h(3 - h)$. Boundary correlators of $\mathcal{O}$ are then related by a rescaling to bulk Φ correlators whose arguments are pushed to the boundary at $\mathcal{I}^+$. As in AdS/CFT , a particular classical bulk wavefunction of Φ can be associated to a boundary insertion of $\mathcal{O}$ (at the linearized level) by symmetries: it must scale with weight h under the isometry corresponding to dilations, obey the lowest-weight condition, and be invariant under rotations around the point of the boundary insertion. The resulting wavefunction is a type of bulk-to-boundary Green function. Interestingly[13], the (lowest) highest-weight modes can also be identified as (anti) quasinormal modes for the static patch of de Sitter, as constructed in [11, 12]. In this section we determine this wavefunction explicitly, regulate the singularities, generalize it to multi-particle insertions and define a symplectic product. In the following section we will then use these classical objects to construct the associated dual bulk quantum states and their inner products.

3.1 Highest and lowest weight wavefunctions

In this subsection we give expressions for the classical wavefunctions, associated to lowest (highest) weight primary operator insertions at the south (north) pole of $\mathcal{I}^+$ in terms of rescaled Green functions in the limit that one argument is pushed to $\mathcal{I}^+$. These wavefunctions each comes in a Neumann and a Dirichlet flavor, denoted $\Phi_{lw}^\pm(x)$ ($\Phi_{hw}^\pm(x)$) depending on whether the weight of the dual operator insertion is h_+ or h_- .

The relevant Green functions are⁴

$$G_\pm(x; x') \equiv G_E(x; x') \pm G_E(x; x'_A) = \frac{1}{8\pi^2} \left(\frac{1}{1 - P(x; x')} \pm \frac{1}{1 + P(x; x')} \right) \quad (3.1)$$

with G_E the Wightman function for the Euclidean vacuum given in equation (2.42). G_- (G_+) obeys Neumann (Dirichlet) boundary conditions at $\mathcal{I}^+$ away from $x = x'$. These are for

⁴Note that $G_+(G_-)$ is even (odd) under the antipodal map. Combinations of the Euclidean Green function with such properties have been previously studied in the context of elliptic Z_2 -identification of de Sitter space [17, 18, 19] which may be related to our construction.

$m^2\ell^2 = 2$ the Green functions with future boundary conditions as discussed in [10]. We have normalized them so that they have the Hadamard form for the short-distance singularity. In the Neumann case we begin with G_- , which is (using the mode decomposition (2.40)) given by

$$G_-(x; x') = \sum (-)^L (\phi_{Lj}^-(x) \phi_{Lj}^-(x') + i \phi_{Lj}^+(x) \phi_{Lj}^-(x')). \quad (3.2)$$

From this we construct the rescaled Green function

$$\Phi_{lw}^-(x; t') = e^{h-t'} G_-(x; t', \Omega_{SP}), \quad (3.3)$$

in which the second argument is placed at the south pole Ω_{SP} where $\psi' = 0$. One may then check that (ignoring singularity prescriptions)

$$\Phi_{lw}^-(x) \equiv \lim_{t' \rightarrow \infty} \Phi_{lw}^-(x; t') = \frac{1}{2\pi^2(\sinh t - \cos \psi \cosh t)}. \quad (3.4)$$

Using (3.2) and the asymptotics (2.11) one finds that near $\mathcal{I}^+$ (not ignoring singularities)

$$\begin{aligned} \Phi_{lw}^-(x) &= 8 \sum (-)^L \left(\frac{e^{-t}}{2L+2} Y_{Lj}(\Omega) Y_{Lj}(\Omega_{SP}) + i e^{-2t} Y_{Lj}(\Omega) Y_{Lj}(\Omega_{SP}) + \mathcal{O}(e^{-3t}) \right) \\ &= 8e^{-t} \Delta_-(\Omega, \Omega_{SP}) + 8i \frac{e^{-2t}}{\sqrt{h}} \delta^3(\Omega - \Omega_{SP}) + \mathcal{O}(e^{-3t}) \end{aligned} \quad (3.5)$$

Let us now confirm that $\Phi_{lw}^-(x)$ has the same symmetries as an insertion of a primary operator $\mathcal{O}(\Omega_{SP})$ at the south pole of $\mathcal{I}^+$. First we note that the choice of a point on $\mathcal{I}^+$ breaks $SO(4, 1)$ to $SO(3) \times SO(1, 1)$. Both $\Phi_{lw}^-(x)$ and $\mathcal{O}(\Omega_{SP})$ are manifestly invariant under the $SO(3)$ spatial rotations. The generator of $SO(1, 1)$ dilations, denoted L_0 , acts on $\mathcal{O}(\Omega_{SP})$ as

$$[L_0, \mathcal{O}(\Omega_{SP})] = h_- \mathcal{O}(\Omega_{SP}). \quad (3.6)$$

In the bulk it is generated by the Killing vector field

$$L_0 = \cos \psi \partial_t - \tanh t \sin \psi \partial_\psi, \quad (3.7)$$

where the south pole is $\psi = 0$. dS invariance implies

$$(L_0 - L'_0) G_-(x, x') = 0. \quad (3.8)$$

It follows from this together with the definition (3.4) that the wavefunction obeys

$$L_0 \Phi_{lw}^-(x) = h_- \Phi_{lw}^-(x). \quad (3.9)$$

By construction it obeys the wave equation

$$(\nabla^2 - m^2) \Phi_{lw}^-(x) = 0. \quad (3.10)$$

Acting on $SO(3)$ invariant symmetric functions we have

$$\ell^2 \nabla^2 = -L_0(L_0 - 3) + M_{-k} M_{+k}, \quad (3.11)$$

where the 6 Killing vector fields $M_{\pm k}$ (given in Appendix A) are the raising and lowering operators for L_0 and we sum over k . It then follows that

$$M_{-k} M_{+k} \Phi_{lw}^-(x) = (m^2 \ell^2 - h_-(3 - h_-)) \Phi_{lw}^-(x) = 0, \quad (3.12)$$

and hence

$$M_{+k} \Phi_{lw}^-(x) = 0. \quad (3.13)$$

which corresponds to the lowest-weight condition for the $\mathcal{O}$

$$[M_{+k}, \mathcal{O}(\Omega_{SP})] = 0. \quad (3.14)$$

It may be shown that these symmetries uniquely determine the solution. Hence Φ_{lw}^- is identified as the classical wavefunction associated to the insertion of the primary $\mathcal{O}$ at the south pole.

A parallel argument leads to the dual of a highest weight operator insertion at the north pole. The wavefunction is

$$\Phi_{hw}^-(x; t') = \lim_{t' \rightarrow \infty} e^{h-t'} G_-(x; t', \Omega_{NP}). \quad (3.15)$$

This obeys the relations

$$M_{-k} \Phi_{hw}^-(x) = 0, \quad L_0 \Phi_{hw}^-(x) = -h_- \Phi_{hw}^-(x), \quad (3.16)$$

and has the asymptotic behavior

$$\Phi_{hw}^-(x) = 8e^{-t}\Delta_-(\Omega, \Omega_{NP}) + 8i\frac{e^{-2t}}{\sqrt{h}}\delta^3(\Omega - \Omega_{NP}) + \mathcal{O}(e^{-3t}). \quad (3.17)$$

Similar formulae apply to the Dirichlet case by beginning with G_+ in the above construction and replacing $+\leftrightarrow -$. For example

$$\Phi_{hw}^+(x) = -8e^{-2t}\Delta_+(\Omega, \Omega_{NP}) - 8i\frac{e^{-t}}{\sqrt{h}}\delta^3(\Omega - \Omega_{NP}) + \mathcal{O}(e^{-3t}). \quad (3.18)$$

We see from the above that the highest-weight wavefunction is smooth on the future horizon of the southern static patch dS_4 , and hence related to the quasinormal modes found in [11, 12]. The lowest quasinormal mode which is invariant under the $SO(3)$ of the static dS_4 is exactly the Φ_{hw}^- with $h_- = 1$ while the second lowest $SO(3)$ -invariant quasinormal mode corresponds to Φ_{hw}^+ with $h_+ = 2$. Lowest weight states are smooth on the past horizon and hence related to anti-quasinormal modes.

3.2 General multi-operator insertions

In the preceding subsections we found the bulk duals of primary operators inserted at the north/south pole in the coordinates (2.3). This can be generalized to insertions at an arbitrary point on $\mathcal{I}^+$ with a general time slicing near $\mathcal{I}^+$. Let us introduce coordinates $x \sim (y^i, t)$ such that near $\mathcal{I}^+$

$$ds_4^2 \rightarrow -dt^2 + e^{2t}h_{ij}(y)dy^i dy^j, \quad i, j = 1, 2, 3. \quad (3.19)$$

The dual wavefunction is then the $t' \rightarrow \infty$ limit of the rescaled Green function, denoted by

$$\Phi_{y_1}^\pm(x) = \lim_{t' \rightarrow \infty} e^{h_\pm t'} G_\pm(x; t', y_1). \quad (3.20)$$

For the special cases of operator insertions at the north or south pole in global coordinates these reduce to our previous expressions. Note that coordinate transformations of the form $t \rightarrow t + f(y)$ induce a conformal transformation on $\mathcal{I}^+$

$$h_{ij} \rightarrow e^{2f(y)}h_{ij}, \quad \Phi_{y_1}^\pm(x) \rightarrow e^{h_\pm f(y)}\Phi_{y_1}^\pm(x), \quad (3.21)$$

as appropriate for a conformal field of weight $h_\pm$. Hence the relative normalization in (3.4) will depend on the conformal frame at $\mathcal{I}^+$.

One may also consider multi-operator insertions such as $\mathcal{O}(y_1)\mathcal{O}(y_2)$ in the CFT_3 at $\mathcal{I}^+$.

At the level of free field theory considered here these are associated to a bilocal wavefunction in the product of two bulk scalar fields

$$\Phi_{y_1}(x_1)\Phi_{y_2}(x_2). \quad (3.22)$$

We will use Φ_Ω to denote these wavefunctions when working in global coordinates (2.3). We note that in such coordinates near $\mathcal{I}^+$ for an insertion at a general point

$$\Phi_{\Omega_1}^\pm(t, \Omega) = \mp 8e^{-h\pm t} \Delta_\pm(\Omega, \Omega_1) \mp 8i \frac{e^{-h\mp t}}{\sqrt{h}} \delta^3(\Omega - \Omega_1) + \mathcal{O}(e^{-3t}). \quad (3.23)$$

3.3 Klein-Gordon inner product

We wish to define an inner product between e.g. two Neumann wavefunctions $\Phi_{\Omega_1}^-$ and $\Phi_{\Omega_2}^-$. Later on we will compare this to the inner product on the CFT_3 Hilbert space and the two-point function of $\mathcal{O}$ on S^3 . One choice is to take a global spacelike S^3 slice in the interior and define the Klein-Gordon inner product

$$\langle \Phi_{\Omega_1}^- | \Phi_{\Omega_2}^- \rangle_{S^3} \equiv i \int_{S^3} d^3 \Sigma^\mu \Phi_{\Omega_1}^{-*} \overleftrightarrow{\partial}_\mu \Phi_{\Omega_2}^-. \quad (3.24)$$

This integral does not depend on the choice of S^3 which can be pushed up to $\mathcal{I}^+$. One may then see immediately from (3.23) that there are two nonzero terms proportional to Δ_- giving

$$\langle \Phi_{\Omega_1}^- | \Phi_{\Omega_2}^- \rangle_{S^3} = 16 \Delta_-(\Omega_1 - \Omega_2) \quad (3.25)$$

One may also define an inner product not on global spacelike S^3 slices, but on a spacelike R^3 slice which ends on an S^2 on $\mathcal{I}^+$. The result is invariant under any deformation of the S^2 which does not cross the insertion point. To be definite, we take the S^2 to be the equator, Ω_1 to be in the northern hemisphere and Ω_2 to be in the southern hemisphere, and the slice to be R_S^3 which intersects the south pole. One then finds, pushing R_S^3 up to the southern hemisphere of $\mathcal{I}^+$

$$\langle \Phi_{\Omega_1}^- | \Phi_{\Omega_2}^- \rangle_{R_S^3} \equiv i \int_{R_S^3} d^3 \Sigma^\mu \Phi_{\Omega_1}^{-*} \overleftrightarrow{\partial}_\mu \Phi_{\Omega_2}^- = 8 \Delta_-(\Omega_1 - \Omega_2). \quad (3.26)$$

Similarly, the inner product between two Dirichlet wavefunctions is given by

$$\langle \Phi_{\Omega_1}^+ | \Phi_{\Omega_2}^+ \rangle_{R_S^3} = -8 \Delta_+(\Omega_1 - \Omega_2). \quad (3.27)$$

4 The southern Hilbert space

We now turn to the issue of bulk quantum states. Quantum states in dS are often discussed, as in section 2, in terms of a Hilbert space built on the global S^3 slices. The structure of the vacua and Green functions for such states was described in section 2. However dS has the unusual feature that there are geodesically complete topologically R^3 spacelike slices which end on an S^2 in $\mathcal{I}^+$, which we will typically take to be the equator. Examples of these are the hyperbolic slices, the quantization on which was studied in detail in [15]. We will see that the quantum states built on these R^3 slices are natural objects in dS/CFT. An S^2 in $\mathcal{I}^+$ is in general the boundary of a “northern” slice, denoted R_N^3 and a “southern” slice denoted R_S^3 . The topological sum obeys $R_S^3 \cup R_N^3 = S^3$. Hence the relation of the southern and northern Hilbert spaces on R_S^3 and R_N^3 to that on S^3 is like that of the left and right Rindler wedges to that of global Minkowski space. It is also like the relation of the Hilbert spaces of the northern and southern causal diamonds to that of global dS. However the diamond Hilbert spaces in dS quantum gravity are problematic in quantum gravity with a fluctuating metric because it is hard to find sensible boundary conditions.

A strong motivation for considering the $R_{S,N}^3$ slices comes from the picture of a state in the boundary CFT₃. The state-operator correspondence in CFT₃ begins with an insertion of a (primary or descendant) operator $\mathcal{O}$ at the south pole of S^3 , and then defines a quantum state as a functional of the boundary conditions on an S^2 surrounding the south pole. For every object in the CFT₃, we expect a holographically dual object in the bulk dS₄ theory. The dual bulk quantum state must somehow depend on the choice of S^2 in $\mathcal{I}^+$. Hence it is natural to define the bulk state on the R^3 slice which ends on this S^2 in $\mathcal{I}^+$. This is how holography works in AdS/CFT: CFT states live on the boundaries of the spacelike slices used to define the bulk states.

4.1 States

In order to define quantum states on R_S^3 , we first note that modes of the scalar field operator $\hat{\Phi}(\Omega, t)$ are labeled by operators $\hat{\Phi}^\pm(\Omega)$ defined on $\mathcal{I}^+$ via the relation

$$\lim_{t \rightarrow \infty} \hat{\Phi}(\Omega, t) = e^{-h_+ t} \hat{\Phi}^+(\Omega) + e^{-h_- t} \hat{\Phi}^-(\Omega). \quad (4.1)$$

They satisfy the following commutation relation

$$\left[\hat{\Phi}^+(\Omega), \hat{\Phi}^-(\Omega') \right] = \frac{8i}{\sqrt{h}} \delta^3(\Omega - \Omega'). \quad (4.2)$$

We may then decompose these $\mathcal{I}^+$ operators as the sum of two terms

$$\hat{\Phi}^\pm(\Omega) = \hat{\Phi}_N^\pm(\Omega) + \hat{\Phi}_S^\pm(\Omega) \quad (4.3)$$

where the first (second) acts only on R_N^3 (R_S^3). Defining the northern and southern Dirichlet and Neumann vacua by

$$\hat{\Phi}_N^\pm |0_N^\pm\rangle = 0, \quad \hat{\Phi}_S^\pm |0_S^\pm\rangle = 0, \quad (4.4)$$

it follows from the decomposition (4.3) that the global vacua have a simple product decomposition⁵

$$|0^\pm\rangle = |0_N^\pm\rangle |0_S^\pm\rangle. \quad (4.5)$$

Excited southern states may then be built by acting on one of these southern vacua with $\hat{\Phi}_S$. We wish to identify these states with those of the CFT₃ on S^2 .

In the higher-spin dS/CFT correspondence there are actually two CFT₃'s living on $\mathcal{I}^+$: the free $Sp(N)$ model, associated to Neumann boundary conditions, and the critical $Sp(N)$ model, associated to Dirichlet boundary conditions. Since the field operators $\hat{\Phi}_S$ acting on $|0_S^+\rangle$ ($|0_S^-\rangle$) obeys, according to equation (2.26), Neumann (Dirichlet) boundary conditions near the southern hemisphere of $\mathcal{I}^+$, it is natural to identify

$$\begin{aligned} |0_S^+\rangle &\sim \text{free } Sp(N) \text{ vacuum} \\ |0_S^-\rangle &\sim \text{critical } Sp(N) \text{ vacuum.} \end{aligned} \quad (4.6)$$

Next we want to consider excited states and their duals. To be specific we consider the Neumann theory built on $|0_S^+\rangle$. Parallel formulae apply to the Dirichlet case. Operator versions of the classical wavefunctions $\Phi_\Omega^-(x)$ are constructed as

$$\hat{\Phi}_{\Omega_S}^- \equiv \left\langle \Phi_{\Omega_S}^- | \hat{\Phi} \right\rangle_{R_S^3}, \quad (4.7)$$

where Ω_S is presumed to lie on the southern hemisphere. We can make a quantum state

$$|\Omega_S^-\rangle \equiv \hat{\Phi}_{\Omega_S}^- |0_S^+\rangle = \hat{\Phi}^-(\Omega_S) |0_S^+\rangle, \quad (4.8)$$

where in the last line we used (3.23). By construction this will be a lowest weight state, and

⁵Of course a general quantum state on S^3 is a sum of products of northern and southern states, and reduces to a southern density matrix, not a pure state, after tracing over the northern Hilbert space. We shall see this explicitly below for the case of the Euclidean vacuum.

we therefore identify it as the bulk dual to the CFT_3 state created by the primary operator $\mathcal{O}$ dual to the field Φ .

This connection leads to an interesting nonperturbative dS exclusion principle [14]. The operator $\mathcal{O}$ has a representation in the $Sp(N)$ theory as

$$\mathcal{O} = \Omega_{AB} \eta^A \eta^B, \quad A, B = 1, \dots, N, \quad (4.9)$$

where η^A are N anticommuting real scalars and Ω_{AB} is the quadratic form on $Sp(N)$. It follows that

$$\mathcal{O}^{\frac{N}{2}+1} = 0. \quad (4.10)$$

Bulk-boundary duality and the state-operator relation described above then implies the nonperturbative relation

$$\left[\hat{\Phi}^\pm(\Omega) \right]^{\frac{N}{2}+1} = 0. \quad (4.11)$$

Hence the quantum field operators $\hat{\Phi}^\pm(\Omega)$ are $\frac{N}{2}$ -adic. One is not allowed to put more than $\frac{N}{2}$ quanta in any given quasinormal mode. This is similar to the stringy exclusion principle for AdS [20] and may be related to the finiteness of dS entropy. Nonperturbative phenomena due to related finite N effects in the $O(N)$ case have been discussed in [21]. We hope to investigate further the consequences of this dS exclusion principle.

4.2 Norm

Having identified the bulk duals of the boundary CFT_3 states, we wish to describe the bulk dual of the CFT_3 norm. The standard bulk norm is defined by $\Phi(x) = \Phi^\dagger(x)$. However this norm is not unique. It has been argued for a variety of reasons beginning in [3] that it is appropriate to modify the norm in the context of dS – see also [16, 22]. Here we have the additional problem that this standard norm is divergent for states of the form (4.8). We now construct the modified norm for states on R_3^S by demanding that it is equivalent to the CFT_3 norm. The construction here generalizes to dS_4 the one given in [16] for dS_3 .

The bulk action of dS Killing vectors $K_A^\mu \partial_\mu$ on a scalar field is generated by the integral over any global S^3 slice

$$\hat{\mathcal{L}}_A = \int_{S^3} d^3 \Sigma^\mu T_{\mu\nu} K_A^\nu, \quad (4.12)$$

where $T_{\mu\nu}$ is the bulk stress tensor constructed from the operator $\hat{\Phi}$. If we take $\hat{\Phi}^\dagger(x) = \hat{\Phi}(x)$, then $\hat{\mathcal{L}}_A = \hat{\mathcal{L}}_A^\dagger$ which is not what we want. The CFT_3 states are in representations of the $SO(3,2)$ conformal group. These arise from analytic continuation of the 10 $SO(4,1)$

conformal Killing vectors on S^3 which are the boundary restrictions of the bulk dS_4 Killing vectors $K_A^\mu \partial_\mu$. Usually, the standard CFT_3 norm has a self-adjoint dilation operator $\mathcal{L}_0$ generating $-i \sin \psi \partial_\psi$ as well as 3 self-adjoint $SO(3)$ rotation operators $\mathcal{J}_k$. The remaining 6 raising and lowering operators $\mathcal{L}_{\pm k}$ arising from the Killing vectors $iM_{\pm k}$ (described in the appendix) then obey $\mathcal{L}_{\pm k}^\dagger = \mathcal{L}_{\mp k}$ in the conventional CFT_3 norm.

To obtain an adjoint with the desired properties, we define the modified adjoint

$$\hat{\Phi}^\dagger(x) = \mathcal{R} \hat{\Phi}(x) \mathcal{R} = \hat{\Phi}(Rx), \quad (4.13)$$

where here and hereafter $\dagger$ denotes the bulk modified adjoint. The reflection operator $\mathcal{R}$ is the discrete isometry of S^3 which reflects through the S^2 equator $R(\psi, \theta, \phi) = (\pi - \psi, \theta, \phi)$ along with complex conjugation. In particular, it maps the south pole to the north pole while keeping the equator invariant. This implies that $\mathcal{L}_0$ (generating iL_0) and $\mathcal{J}_k$ are self adjoint while

$$\mathcal{L}_{\pm k}^\dagger = -i \int_{S^3} d\Sigma^\mu(x) T_{\mu\nu}(Rx) M_{\pm k}^\nu(x) = -i \int_{S^3} d\Sigma^\mu(x) T_{\mu\nu}(x) M_{\pm k}^\nu(Rx) = \mathcal{L}_{\mp k}. \quad (4.14)$$

Hence we have constructed an adjoint admitting the desired $SO(3,2)$ action. We do not know whether or not it is unique.

The action of $\mathcal{R}$ maps an operator defined on the southern hemisphere to one defined on the southern hemisphere of $\mathcal{I}^+$ according to

$$\hat{\Phi}^{\pm\dagger}(\Omega) = \hat{\Phi}^\pm(\Omega_R), \quad (4.15)$$

Hence the action of $\mathcal{R}$ exchanges the northern and southern hemispheres, and maps a southern $\mathcal{I}^+$ state to a northern one. Therefore it cannot on its own define an adjoint within the southern Hilbert space. For this we must combine (4.13) with a map from the north to the south. Such a map is provided by the Euclidean vacuum. The global Euclidean bra state (constructed with the standard adjoint) can be decomposed in terms of a basis of northern and southern bra states

$$\langle 0_E | = \sum_{m,n} E_{mn} \langle m_S | \langle n_N |. \quad (4.16)$$

We then define the modified adjoint of an arbitrary southern state $|\Psi_S\rangle$ by

$$|\Psi_S\rangle^\dagger \equiv \langle 0_E | \mathcal{R} | \Psi_S \rangle. \quad (4.17)$$

We will denote the corresponding inner product by an S subscript

$$\langle \Psi'_S | \Psi_S \rangle_S \equiv (|\Psi'_S\rangle^\dagger) | \Psi_S \rangle. \quad (4.18)$$

For example choosing the basis so that

$$\mathcal{R} | m_S \rangle = | m_N \rangle \quad (4.19)$$

we have

$$\langle m_S | n_S \rangle_S = E_{nm}. \quad (4.20)$$

In particular one finds

$$\langle 0_S^+ | 0_S^+ \rangle_S = \langle 0_E | (| 0_S^+ \rangle \mathcal{R} | 0_S^+ \rangle) = 1. \quad (4.21)$$

Let us now compute the norm of the southern state $|\Omega_S^- \rangle$ in (4.8). The action of $\mathcal{R}$ gives a northern state which we will denote $|R\Omega_S^- \rangle$. The norm is then

$$\langle \Omega_S^- | \Omega_S^- \rangle_S = \langle 0_E | (| \Omega_S^- \rangle | R\Omega_S^- \rangle) = \langle 0_E | \hat{\Phi}^-(\Omega_S) \hat{\Phi}^-(R\Omega_S) | 0^+ \rangle. \quad (4.22)$$

Using the relation

$$| 0_E \rangle = N_0 e^{-\frac{1}{16} \int d^3\Omega d^3\Omega' \hat{\Phi}^+(\Omega) \Delta_-(\Omega, \Omega') \hat{\Phi}^+(\Omega')} | 0^- \rangle \quad (4.23)$$

we find

$$\langle \Omega_S^- | \Omega_S^- \rangle_S = 8\Delta_-(\Omega_S, R\Omega_S). \quad (4.24)$$

This is proportional to the S^3 two-point function of a dimension h_- primary at the points Ω_S and $R\Omega_S$. The analogous computation in the Dirichlet theory gives

$$\langle \Omega_S^+ | \Omega_S^+ \rangle_S = -8\Delta_+(\Omega_S, R\Omega_S). \quad (4.25)$$

5 Boundary dual of the bulk Euclidean vacuum

In the preceding section we have argued that dS/CFT maps CFT₃ states on an S^2 in $\mathcal{I}^+$ to bulk states on the southern slice ending on the S^2 . A generic state in a global dS slice does not restrict to a pure southern state. However we can always define a density matrix by tracing over the northern Hilbert space. In particular, such a southern density matrix ρ_S^E can be associated to the global Euclidean vacuum $| 0_E \rangle$. The choice of an equatorial S^2 in $\mathcal{I}^+$ breaks the $SO(4, 1)$ symmetry group down to $SO(3, 1)$, which also preserves the hyperbolic

slices ending on the S^2 . ρ_E^S must be invariant under this $SO(3, 1)$. In fact ρ_E^S follows from results in [15]. Writing the quadratic Casimir of $SO(3, 1)$ as $C_2 = -(1 + p^2)$, it was shown, in a basis which diagonalizes p , that

$$\rho_E^S = N_1 e^{-2\pi p}, \quad (5.1)$$

where N_1 is determined by $\text{Tr} \rho_E^S = 1$. It would be interesting to investigate this further and compute the entropy $S = -\text{Tr} \rho_E^S \ln \rho_E^S$ in the $Sp(N)$ model.

6 Pseudounitariness and the C-norm in the $Sp(N)$ CFT₃

In this section we consider the $Sp(N)$ model (where N is even) and compare the norms to those computed above. The action is

$$I_{Sp(N)} = \frac{1}{8\pi} \int d^3x \left[\delta^{ij} \delta_{ab} \partial_i \bar{\chi}^a \partial_j \chi^b + m^2 \bar{\chi} \chi + \lambda (\bar{\chi} \chi)^2 \right], \quad (6.1)$$

where $\chi^a (a = 1, \dots, \frac{N}{2})$ is a complex anticommuting scalar and $\bar{\chi} \chi \equiv \delta_{ab} \bar{\chi}^a \chi^b$. This has a global $Sp(N)$ symmetry and we restrict to $Sp(N)$ singlet operators.⁶ For the free theory $m = \lambda = 0$ while the critical theory is obtained by flowing to a nontrivial fixed point λ_F . The $Sp(N)$ theory is not unitary in the sense that in the standard norm following from (6.1) one has that [9]

$$H \neq H^\dagger \quad (6.2)$$

and $\langle \Psi' | \Psi \rangle$ is not preserved. Nevertheless, as detailed in [9], there exists an operator C with the properties

$$C^\dagger C = C^2 = 1, \quad C \chi^\dagger C = \bar{\chi}, \quad C H^\dagger C = H, \quad C |0\rangle = |0\rangle. \quad (6.3)$$

To write it in real fields, for e.g., in the case of $Sp(2)$, writing the real and imaginary part of χ as η_1 and η_2 , the action of C becomes $\eta_2 = C \eta_1^\dagger C$. One may then define a “pseudounitary” C -inner product

$$\langle \Psi' | \Psi \rangle_C \equiv \langle \Psi' | C | \Psi \rangle \quad (6.4)$$

which is preserved under hamiltonian time evolution. Such hamiltonians are pseudohermitian and are similar to those studied in [23]. We note that the norm is not positive definite.

⁶The $U(\frac{N}{2})$ theory has the same action but is restricted only to $U(\frac{N}{2})$ singlets.

Inserting an operator O_i constructed from χ^a at the south pole gives a functional of the boundary conditions on the equatorial S^2 which we define as the state $|O_i\rangle$. This is the standard state-operator correspondence. An inner product for such states associated to O_i and O_j can be defined by the two point function with O_i at one pole and O_j at the other. It follows from (6.3) that this is the C -inner product for the states $|O_i\rangle$ and $|O_j\rangle$:

$$\langle O_i|O_j\rangle_C = \langle O_i|C|O_j\rangle = \langle O_i^\dagger C O_j\rangle = \langle O_i O_j\rangle. \quad (6.5)$$

In the last line, we used the fact that the (singlet) currents in the $Sp(N)$ models satisfy $CO_i^\dagger C = O_i$ since $C(\bar{\chi}\chi)^\dagger C = \bar{\chi}\chi$. For primary operators of weight h_i we then have [6]

$$\langle O_i|O_i\rangle_C = -N\Delta_{h_i}(\Omega_{NP}, \Omega_{SP}). \quad (6.6)$$

Hence it is the C-norm which maps under the state-operator correspondence to the Zamolodchikov norm defined as the Euclidean two point function on S^3 . As seen in [6] this C-norm then agrees with the bulk inner product (4.24)-(4.25) of the dual state for the scalar case.⁷ Moreover, as the bulk and CFT₃ norms assign the same hermiticity properties to the $SO(4,1)$ generators, this result will carry over to descendants of the primaries. A generalization of this construction to all spins seems possible.

One of the puzzling features of dS/CFT is that the dual CFT cannot be unitary in the ordinary sense. This is not a contradiction of any kind because unitarity of the Euclidean CFT is not directly connected to any spacetime conservation law. At the same time quantum gravity in dS – and its holographic dual – should have some good property replacing unitarity in the AdS case. It is not clear what that good property is. The appearance of a pseudounitary structure in the case of dS/CFT analyzed here is perhaps relevant in this regard.

Acknowledgements

It has been a great pleasure discussing this work with Dionysios Anninos, Daniel Harlow, Tom Hartman, Daniel Jafferis, Matt Kleban and Steve Shenker. This work was supported in part by DOE grant DE-FG02-91ER40654 and the Fundamental Laws Initiative at Harvard.

⁷A factor of i^h explained in [6] relating operator insertions to bulk fields makes the two-point function (4.24) negative.

A Appendix: dS₄ Killing vectors

The 10 Killing vectors of dS₄ are given by:

$$\begin{aligned}
L_0 &= \cos \psi \partial_t - \tanh t \sin \psi \partial_\psi \\
M_{\mp 1} &= \pm \sin \psi \sin \theta \sin \phi \partial_t + (1 \pm \tanh t \cos \psi) \sin \theta \sin \phi \partial_\psi \\
&\quad + (\cot \psi \pm \tanh t \csc \psi) (\cos \theta \sin \phi \partial_\theta + \csc \theta \cos \phi \partial_\phi) \\
M_{\mp 2} &= \pm \sin \psi \sin \theta \cos \phi \partial_t + (1 \pm \tanh t \cos \psi) \sin \theta \cos \phi \partial_\psi \\
&\quad + (\cot \psi \pm \tanh t \csc \psi) (\cos \theta \cos \phi \partial_\theta - \csc \theta \sin \phi \partial_\phi) \\
M_{\mp 3} &= \pm \sin \psi \cos \theta \partial_t + (1 \pm \tanh t \cos \psi) \cos \theta \partial_\psi - (\cot \psi \pm \tanh t \csc \psi) \sin \theta \partial_\theta \\
J_1 &= \cos \phi \partial_\theta - \sin \phi \cot \theta \partial_\phi \\
J_2 &= -\sin \phi \partial_\theta - \cos \phi \cot \theta \partial_\phi \\
J_3 &= \partial_\phi.
\end{aligned} \tag{A.1}$$

For each k , the $M_{\pm k}$ and L_0 form a $SO(2, 1)$ subalgebra satisfying

$$[M_{+k}, M_{-k}] = 2L_0, \quad [L_0, M_{+k}] = -M_{+k}, \quad [L_0, M_{-k}] = M_{-k}. \tag{A.2}$$

As mentioned in the text, acting on $SO(3)$ -invariant functions, we have

$$\ell^2 \nabla^2 = -L_0(L_0 - 3) + M_{-k}M_{+k}, \tag{A.3}$$

where k is summed over $k = 1, 2, 3$.

The conformal Killing vectors of the S^3 are given by the restriction of dS₄ Killing vectors on $\mathcal{I}^+$:

$$\begin{aligned}
L_0 &= -\sin \psi \partial_\psi \\
M_{\mp 1} &= (1 \pm \cos \psi) \sin \theta \sin \phi \partial_\psi + (\cot \psi \pm \csc \psi) (\cos \theta \sin \phi \partial_\theta + \csc \theta \cos \phi \partial_\phi) \\
M_{\mp 2} &= (1 \pm \cos \psi) \sin \theta \cos \phi \partial_\psi + (\cot \psi \pm \csc \psi) (\cos \theta \cos \phi \partial_\theta - \csc \theta \sin \phi \partial_\phi) \\
M_{\mp 3} &= (1 \pm \cos \psi) \cos \theta \partial_\psi - (\cot \psi \pm \csc \psi) \sin \theta \partial_\theta. \\
J_1 &= \cos \phi \partial_\theta - \sin \phi \cot \theta \partial_\phi \\
J_2 &= -\sin \phi \partial_\theta - \cos \phi \cot \theta \partial_\phi \\
J_3 &= \partial_\phi.
\end{aligned} \tag{A.4}$$

References

- [1] A. Strominger, “The dS/CFT correspondence,” JHEP **0110**, 034 (2001) [arXiv:hep-th/0106113]; “Inflation and the dS/CFT correspondence,” JHEP **0111**, 049 (2001) [arXiv:hep-th/0110087].
- [2] C. M. Hull, “Timelike T duality, de Sitter space, large N gauge theories and topological field theory,” JHEP **9807**, 021 (1998) [hep-th/9806146].
- [3] E. Witten, “Quantum gravity in de Sitter space,” hep-th/0106109.
- [4] J. M. Maldacena, “Non-Gaussian features of primordial fluctuations in single field inflationary models,” JHEP **0305**, 013 (2003) [astro-ph/0210603].
- [5] D. Harlow and D. Stanford, “Operator Dictionaries and Wave Functions in AdS/CFT and dS/CFT,” arXiv:1104.2621 [hep-th].
- [6] D. Anninos, T. Hartman and A. Strominger, “Higher Spin Realization of the dS/CFT Correspondence,” arXiv:1108.5735 [hep-th].
- [7] M. A. Vasiliev, “Consistent equation for interacting gauge fields of all spins in (3+1)-dimensions,” Phys. Lett. B **243**, 378 (1990).
- [8] M. A. Vasiliev, “Higher spin gauge theories: Star-product and AdS space,” arXiv:hep-th/9910096.
- [9] A. LeClair, M. Neubert, “Semi-Lorentz invariance, unitarity, and critical exponents of symplectic fermion models,” JHEP **0710**, 027 (2007). [arXiv:0705.4657 [hep-th]].
- [10] D. Anninos, G. S. Ng and A. Strominger, “Future Boundary Conditions in De Sitter Space,” JHEP **1202**, 032 (2012) [arXiv:1106.1175 [hep-th]].
- [11] A. Lopez-Ortega, “Quasinormal modes of D-dimensional de Sitter spacetime,” Gen. Rel. Grav. **38**, 1565 (2006) [gr-qc/0605027].
- [12] D. Anninos, S. A. Hartnoll and D. M. Hofman, “Static Patch Solipsism: Conformal Symmetry of the de Sitter Worldline,” Class. Quant. Grav. **29**, 075002 (2012) [arXiv:1109.4942 [hep-th]].
- [13] D. Anninos, private communication.

- [14] M. Kleban, G. S. Ng, S. Shenker and A. Strominger, unpublished and in progress.
- [15] M. Sasaki, T. Tanaka and K. Yamamoto, “Euclidean vacuum mode functions for a scalar field on open de Sitter space,” *Phys. Rev. D* **51**, 2979 (1995) [gr-qc/9412025].
- [16] R. Bousso, A. Maloney, A. Strominger, *Phys. Rev.* **D65**, 104039 (2002). [hep-th/0112218].
- [17] N. G. Sanchez, “Quantum Field Theory And The ‘Elliptic Interpretation’ Of De Sitter Space-time,” *Nucl. Phys. B* **294**, 1111 (1987).
- [18] G. W. Gibbons, “The Elliptic Interpretation Of Black Holes And Quantum Mechanics,” *Nucl. Phys. B* **271**, 497 (1986).
- [19] M. K. Parikh, I. Savonije and E. P. Verlinde, “Elliptic de Sitter space: dS/Z(2),” *Phys. Rev. D* **67**, 064005 (2003) [hep-th/0209120].
- [20] J. M. Maldacena and A. Strominger, “AdS(3) black holes and a stringy exclusion principle,” *JHEP* **9812**, 005 (1998) [hep-th/9804085].
- [21] S. H. Shenker and X. Yin, “Vector Models in the Singlet Sector at Finite Temperature,” arXiv:1109.3519 [hep-th].
- [22] M. K. Parikh and E. P. Verlinde, “De Sitter holography with a finite number of states,” *JHEP* **0501**, 054 (2005) [hep-th/0410227].
- [23] C. M. Bender, *Repts. Prog. Phys.*, 70, 947 (2007).